

Protection of the Liver by Ischemic Preconditioning: A Review of Mechanisms and Clinical Applications

Rahul S. Koti Alexander M. Seifalian Brian R. Davidson

University Department of Surgery and Liver Transplantation Unit, Royal Free and University College Medical School, University College London and Royal Free Hospital NHS Trust, London, UK

Key Words

Liver · Preconditioning · Ischemia · Reperfusion · Liver injury · Hepatoprotection · Liver transplantation · Liver resection · Nitric oxide · Adenosine · Heat shock proteins · Protein kinase C

Abstract

Ischemic preconditioning refers to the endogenous mechanism of protection against a sustained ischemic insult following an initial, brief ischemic stimulus. Ischemia-reperfusion injury of the liver is a major cause of morbidity and mortality in liver surgery and transplantation and ischemic preconditioning is a promising strategy for improving the outcome of liver surgery. The preconditioning phenomenon was first described in a canine model of myocardial ischemia-reperfusion injury in 1986 and since then has been shown to exist in other organs including skeletal muscle, brain, kidneys, retina and liver. In the liver, the preconditioning effect has been demonstrated in rodents and a recent study has demonstrated human clinical benefits of preconditioning during hemihepatectomies. Ischemic preconditioning has been described as an adaptive response and although the precise mechanism of hepatoprotection from preconditioning is unknown it is likely to be a receptor-mediated process. Several hypotheses have been proposed and this

review assesses possible mechanisms of ischemic preconditioning and its role in hepatic surgery and liver transplantation. The future lies in defining the mechanisms of the ischemic preconditioning effect to allow drug targeting to induce the preconditioning response.

Copyright © 2003 S. Karger AG, Basel

Introduction

Ischemia-reperfusion injury is a major cause of morbidity and mortality following liver surgery and transplantation. Ischemia-reperfusion injury after prolonged ischemia has been shown to occur in virtually all organ systems. Ischemic (and reperfusion) injury to the liver occurs during liver resections performed under temporary inflow occlusion (Pringle manoeuvre) or inflow and outflow occlusion commonly used to reduce intraoperative blood loss, and during storage and implantation of livers for transplantation. The liver tolerates prolonged ischemia poorly and safe ischemic times particularly for the diseased liver are not known. Both warm and cold ischemias result in significant liver injury [1], and ischemia reperfusion injury of the liver can result in multiple system organ failure and systemic inflammatory response syndrome (fig. 1).

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2003 S. Karger AG, Basel
0253-4886/03/0205-0383\$19.50/0

Accessible online at:
www.karger.com/dsu

Brian R. Davidson, Professor of Surgery and Liver Transplantation
University Department of Surgery and Liver Transplantation Unit
Royal Free and University College Medical School
Royal Free Hospital, Pond Street, London NW3 2QG (UK)
Tel. +44 207 830 2757, Fax +44 207 830 2688, E-Mail b.davidson@rfc.ucl.ac.uk




Fig. 1. Schematic illustration of pathophysiological events following reperfusion of ischemic liver. ROS = Reactive oxygen species; IL = interleukin; PAF = platelet-activating factor; SIRS = systemic inflammatory response syndrome.

Brief episodes of ischemia followed by a period of reperfusion called ischemic preconditioning (IPC) have been shown to protect organs against subsequent sustained ischemia. IPC was first described by Murry et al. [2] in 1986. In a canine model, they demonstrated that multiple brief ischemic episodes protected the heart from a subsequent sustained ischemic insult. Since then myocardial IPC has been shown to occur in many animal species [3] and in humans [4]. Subsequently, IPC has been demonstrated in other organ systems including skeletal muscle [5], brain [6], spinal cord [7], kidney [8], intestine [9] and liver [10]. Although these studies suggested a preconditioning response in most organ systems the mechanism of the preconditioning effect remains uncertain.

IPC has been described as an endogenous adaptive mechanism for prevention of injury resulting from ischemia reperfusion [2]. The phenomenon is fascinating, as it is easily reproducible and potentially readily applicable in clinical situations. In the liver the preconditioning effect

is a promising strategy in assisting preservation of the liver in clinical situations of anticipated hepatic ischemia such as transplantation and during resection for tumors using hepatic vascular occlusion. Increase in ischemic hepatic tissue tolerance may protect against ischemia-reperfusion injury and assist preservation of livers for transplantation as well improve outcome after surgical procedures. Clearly, identifying the mechanism of preconditioning may allow recognition of a pharmacological agent, to protect the liver from ischemic injury.

The mechanisms underlying the preconditioning effect have not been defined. In contrast, various potential mediators have been proposed and investigated. Most of the data on mediators of preconditioning in organs including the liver has been extrapolated from information gathered in the heart. This article reviews the major developments in characterization of mechanisms of IPC in the liver. In addition, clinical applications of IPC to minimize ischemic injury to the liver have been discussed.

Methods of Search

All the studies were identified by PubMed, ISIS and CAS searches between the years 1966 and September 2002 with the following key words: ischemia, ischemia-reperfusion injury, preconditioning, IPC, hepatoprotection. Other sources include review articles and textbooks.

Evidence that IPC Occurs in the Liver

Studies on Warm Ischemia

Over the last decade many investigators have studied the effects of IPC on regional and global ischemia in the liver, and the evidence is encouraging (see table 1). Lloris-Carsi et al. [10] in 1993 first demonstrated in the rat liver that a single episode of preconditioning with 5 min portal triad clamping followed by 10 min reperfusion showed improved survival and decreased liver enzyme levels after subsequent 90 min ischemia. Hardy et al. [11] showed improved survival in rats undergoing liver resection during 45 min ischemia after prior 5 min ischemia with 10 min reperfusion. Similarly Yoshizumi et al. [12] have demonstrated improved survival and increased tissue ATP with preconditioning in a rat liver resection model. Subsequently the IPC effect in the liver has been reproduced in several in vivo rodent models of partial and global liver ischemia [13–38]. These studies have demonstrated that liver IPC for warm ischemia resulted in

Table 1. Current data on hepatic IPC: published studies

Study group	Year	Species	IPC time min	Ischemia time min	Reperfusion time, min	Hepatic ischemia	Pharmacological manipulations	Parameters assessed	Outcome of IPC	Proposed mechanism
Lloris-Carsi et al. [10]	1993	Rat	1 × 3 (5I + 10R)	90	3 days	Total	Nil	LFTs and survival	70% 3 day survival + ↓ liver enzyme levels	Not addressed
Hardy et al. [11]	1996	Rat	5I + 10R	0–45	1–8 days	Partial	Nil	LFTs, histology and survival	90% 1 day survival, ↑ prothrombin time	Not addressed
Peralta et al. [13]	1996	Rat	10I + 10R	90	90	Partial	Spermine NONOate, L-NAME/bosentan	LFTs, tissue endothelin and NOS activity, histology	↓ GPT and endothelin ↑ cNOS	NO
Kume et al. [29]	1996	Rat	15I	30	10 and 40	Total	Hyperthermia: 42 °C for 15 min	ATP, transaminase, LDH, HSP72, survival	100% survival, ↑ ATP, ↓ transaminase and LDH	HSP72
Peralta et al. [17]	1997	Rat	10I + 10R	90	90	Partial	Spermine NONOate; adenosine; L-NAME	LFTs, HM	↓ transaminases and LDH	Adenosine and NO
Peralta et al. [18]	1998	Rat	2–30I + 10R	90	90	Partial	Adenosine deaminase/xanthine; spermine NONOate	LFTs	↓ transaminases	Adenosine and NO
Yoshizumi et al. [12]	1998	Rat	5I + 10R	40	120	Partial	Nil	LFTs, bile flow, tissue ATP, histology	↓ transaminase, LDH and tissue necrosis, ↑ ATP	Not addressed
Yin et al. [26]	1998	Rat	5I, 10I or 20I + 10R	16–24 h ^a	60 min to 5 days	Cold storage	L-arginine/adenosine; L-NAME	LFTs, bile flow, TNF-α, survival	87.5% 1 day and 75% 5 days graft survival	NO
Adam et al. [57]	1998	Rat	5 or 10I + 10R	24 h ^a	180	Cold storage (UW)	Nil	Bile, transaminases, LDH release, vascular resistance	↑ transaminases, LDH and vascular resistance	Not addressed
Peralta et al. [19]	1999	Rat	10I + 10R	90	90	Partial	Adenosine; adenosine deaminase; DPCPX; DMPX	Transaminases, hepatic perfusion, nitrite/nitrates	Adenosine A2 receptor antagonist abolished	Adenosine A2 receptors and NO
Peralta et al. [20]	1999	Rat	10I + 10R	90	90	Partial	Gadolinium chloride, TNF, L-NAME, spermine NONOate	TNF, transaminases, vascular permeability, edema, MPO, histology	↓ TNF and tissue injury	NO
Yadav et al. [27]	1999	Mouse	10I + 10R	75–90	60–180	Partial and total	Nil	LFTs, hepatocellular apoptosis, survival	↓ apoptosis of hepatocytes and SEC	Modulation of apoptosis cascade
Nakayama et al. [28]	1999	Rat	10I + 10R	45	40 min to 24 h	Total	AdoR A1, A2 agonists and antagonists	LFTs, tissue ATP, histology, survival	↑ adenosine ↓ tissue damage	Adenosine A2 receptors
Zapletal et al. [40]	1999	Rat	5I + 10R	70	30	Partial	Nil	Intravital microscopy	↑ perfusion parameters, ↓ leukocyte adherence	
Nilsson et al. [24]	2000	Rat	10I + 15R	60	60	Total	Dipyridamole	LFTs, HM	↑ peripheral liver blood flow and ↓ transaminase	Adenosine
Howell et al. [25]	2000	Mouse	5I + 10R	30	30 min to 24 h	Partial	Dipyridamole	LFTs, leukocyte/endothelial cell adhesion	↓ endothelial/leukocyte interaction and transaminase	Adenosine
Clavien et al. [43]	2000	Human	10I + 10R	30	30	Total	Nil	LFTs, hepatocellular apoptosis	↓ transaminases and apoptotic sinusoidal lining cells	Modulation of apoptosis cascade
Peralta et al. [21]	2000	Rat	10I + 10R	10–90	90	Partial	SQ-22536, forskolin	Adenine nucleotides, glycogen, glucose-6-P, fructose-6-P, transaminases	Preserved energy metabolism during sustained ischemia	cAMP dependent PKC
Tsumaya et al. [31]	2000	Mice	10/15/20I + 20R	70	1–48 h	Total	Nil	TNF, IL-6, transaminase, histology, survival	↑ survival, ↓ transaminase, TNF, IL-6 and liver necrosis	Not addressed
Peralta et al. [22]	2001	Rat	10I + 10R	90	0–360	Partial	AICAR/8-bromo-AMP/araA; ZVAD; spermine NONOate/ L-NAME	AMPK activity, nucleotides, lactate, transaminases, apoptosis, histology	AMPK activation, ↑ ATP, ↓ lactate and hepatic injury	AMP via PKC
Peralta et al. [23]	2001	Rat	10I + 10R	90	90	Partial	ICAM, P-selectin and TNF blockade; TNF/gadolinium	MPO and lipid peroxidation, vascular permeability, TNF, transaminases, hepatic perfusion, histology, ICAM and P-selectin expression	Remote organ protection by hepatic preconditioning	TNF and P-selectin
Schulz et al. [58]	2001	Pig	10I + 10R × 3	120–200	480 or 300	Total	Nil	ICG clearance, bile flow, transaminases, ATP, glycogen and lactate contents, histology	No protection against prolonged ischemia	Not addressed



Table 1 (continued)

Study group	Year	Species	IPC time min	Ischemia time min	Reperfusion time, min	Hepatic ischemia	Pharmacological manipulations	Parameters assessed	Outcome of IPC	Proposed mechanism
Arai et al. [51]	2001	Rat	5 or 10I + 10R	30 h ^a	15 or 240	Cold storage	Nil	SEC injury, superoxide formation in Kupffer cells, graft survival and TNF after OLT	↑ graft survival and protection contralateral liver	Not addressed
Ricciardi et al. [52]	2001	Pig	15I + 15R	120 ^a	240	Cold storage	PKC inhibitor chelerythrine	Graft function and circulation, LDH, endothelial cell damage, PKC levels	↑ graft function	PKC
Ricciardi et al. [53]	2001	Pig	15I + 15R	120 ^a	240	Cold storage	Tyrosine kinase inhibitor, genistein	graft function and circulation, Tyrosine kinase activity	↑ graft function	Tyrosine kinases
Saito et al. [33]	2001	Rat	10I + 10R	40	6–48 h	Partial	Nil	Transaminases, endothelial cell injury, apoptosis, transcription of IEG's	↓ transaminases, endothelial cell injury, necrosis, apoptosis and IEG transcription	Not addressed
Yamada et al. [32]	2001	Rat	10I + 10R	40–120	Up to 7 days	Partial	Nil	Transaminases, LDH, necrosis, hepatocyte regeneration	↓ transaminases, LDH and necrosis	Not addressed
Zhang et al. [34]	2001	Mice					Nil	Transaminase, LDH, LPO, SOD	↓ transaminases, LDH, LPO and ↑ SOD	Not addressed
Ishii et al. [39]	2001	Rat	10I + 10R	40	6–48 h		Nil	Transaminases, LDH, necrosis, apoptosis, IEG transcription alterations	↓ transaminases, LDH, necrosis, apoptosis, IEG transcription alterations	Not addressed
Peralta et al. [35]	2002	Rat	10I + 10R	90	90	Partial	Xanthine, xanthine oxidase, allopurinol, GSH ester	Xanthine, glutathione, superoxide dismutase, lipid peroxidation, transaminases	↓ xanthine, XOD in liver with ↓ neutrophil accumulation, oxidative stress, and microvascular disorders in lung	Xanthine/XOD pathway for ROS generation
Sindram et al. [50]	2002	Rat	10I + 15R	30 h ^a	60	Cold storage (UW)	N-acetyl-cysteine	SEC detachment, apoptosis, peroxide, gelatinolytic and gelatinase activity	↓ SEC detachment and activities of matrix metalloproteinase	Oxygen free radicals
Rudiger et al. [36]	2002	Mice	10I + 15R	75-120	180	Partial	Nil	Transaminase, apoptosis markers, histology, survival	↓ transaminase, no apoptosis or necrosis, 100% survival for ischemic period up to 75 min but not 120 min	Modulation of apoptosis cascade
Ajamieh et al. [37]	2002	Rat	10I + 10R	90	90	Partial	Ozone	Transaminases, 5'-NT, oxidative stress, histology	↓ hepatocellular injury and oxidative stress	Not addressed
Peralta et al. [14]	2002	Mice	10I + 15R	90	6–24 h	Partial	Gadolinium chloride, TNF, MIP-2, antibodies against TNF and MIP-2	Transaminases, TNF, MIP-2, MDA, MPO, P-selectin expression	Preconditioning and Bcl-2 overexpression together abolished liver injury	Via TNF and MIP-2 inhibition
Serafin et al. [47]	2002	Rat	10I + 10R, 10I + 15R or 5I + 10R	60	2–24 h	Partial	NO donors and inhibitors, glutathione ester	Microcirculation, neutrophil activity, lipid peroxidation	↓ hepatic injury in normal and fatty livers	NO
Teoh et al. [15]	2002	Mice	2–20I + 10R	90	24 h	Partial	Nil	Transaminases, histology,	↓ hepatocellular injury	NF-κB and SAPKs
Fernandez et al. [16]	2002	Rat	10I + 10R	90	16 h	Total	Xanthine, XOD	Transaminases, ROS	↓ liver and lung injury	Via xanthine/XOD blockade
Koti et al. [42]	2002	Rat	5I + 10R	45	120	Partial	L-arginine, L-NAME	Transaminases, NOx, hepatic oxygenation	↓ hepatocellular injury, ↑ intracellular oxygenation	NO
Koti et al. [41]	2002	Rat	5I + 10R	45	120	Partial	L-arginine, L-NAME	Transaminases, NOx, cGMP, microcirculation	↓ hepatocellular injury, ↑ microcirculation	NO
Iwasaki et al. [38]	2002	Rat	10I + 10R	15 × 3 or 45	Up to 180 min	Total	Nil	Transaminases, TNF, histology	↑ protective effect for intermittent than continuous I	Not addressed
Funaki et al. [62]	2002	Mice	15I + 20R	70	0–24 h	Total	Nil	NF-κB activity	↓ NF-κB activation	NF-κB
Ricciardi et al. [63]	2002	Pig	15I + 15R			Cold storage	Genistein, chelerythrine	TK, PKC, NF-κB	TK, PKC, NF-κB activation	PKC and TK, NF-κB

Table 1 (continued)

Study group	Year	Species	IPC time min	Ischemia time min	Reperfusion time, min	Hepatic ischemia	Pharmacological manipulations	Parameters assessed	Outcome of IPC	Proposed mechanism
Arai et al. [48]	1999	Rat	5I + 5R	30 h ^a	15	Cold storage (UW)	Nil	SEC killing, Kupffer cell activation	↓ SEC death and KC activation	Not addressed
Arai et al. [49]	2000	Rat	5I + 5R	30 h ^a		Cold storage (UW)	AdoR A1, A2 agonists and antagonists	SEC killing, SEC cAMP	↓ SEC death and ↑ cAMP	Adenosine A2 receptors via cAMP
Carini et al. [54]	2000	Rat	10I + 10R	90		Hypoxia	PKC stimulators and inhibitors	Hepatocyte viability, pH, Na ⁺ , ATP	↓ hepatocyte cell death	PKC
Carini et al. [55]	2001	Rat	10I + 10R	90		Hypoxia	AdoR A1, A2 agonists and antagonists; PKC, MEK inhibitors	Cell viability, PKC isoenzyme activity, P38 MAPK activity	hepatocyte killing ↓ reduced by 35%	Adenosine A2 receptors, Gi proteins, phospholipase C, PKC, P38 MAPK
Compagnon et al. [56]	2002	Rat	10 anoxia + 10 reoxy-generation	30 + 24 -48 h ^a	60	Warm ischaemia + cold storage	Nil	LDH, ATP, oxygen uptake, protein synthesis	↑ hepatocyte viability, ↑ ATP and protein synthesis	Not addressed

I = Ischemia; R = reperfusion; LFTs = liver function tests; NOS = nitric oxide synthase; cNOS = constitutive nitric oxide synthase; GPT = glutamate pyruvic transaminase; ATP = adenosine triphosphate; LDH = lactate dehydrogenase; HM = hepatic microcirculation; MPO = myeloperoxidase; IL-6 = interleukin-6; AMPK = adenosine monophosphate-activated protein kinase; ICAM = intercellular adhesion molecule; ICG = indocyanine green; OLT = orthotopic liver transplantation; IEGs = immediate early genes; LPO = lipid peroxidase; SOD = superoxide dismutase; XOD = xanthine oxidase; 5'-NT = 5'-nucleotidase; MIP-2 = macrophage inflammatory protein-2; MDA = malondialdehyde; ROS = reactive oxygen species; cAMP = cyclic adenosine monophosphate; SAPKs = stress-activated protein kinases; p38 MAPK = p38 mitogen-activated protein kinase; ↑ = increased; ↓ = decreased.

^a Cold ischemia.

decreased hepatocellular injury [17, 18], increased tissue ATP [12, 22], decreased tumor necrosis factor- α (TNF- α) [19, 31] and IL-6 [31] release, decreased leukocyte/endothelial cell interactions [25], decreased endothelial cell injury [39], increased peripheral liver blood flow [24], increased microcirculation [40, 41], decreased hepatocellular apoptosis [27], preserved energy metabolism [21], increased hepatic intracellular oxygenation [42] and remote organ protection [23]. These studies provide considerable evidence that preconditioning ameliorates ischemia-reperfusion-induced liver injury in the rodent liver. Encouragingly, recent investigations by Clavien et al. [43] have shown that IPC exists in the human liver. In this study, patients undergoing hemihepatectomies under inflow occlusion showed inhibition of sinusoidal endothelial cell (SEC) apoptosis within 30 min of reperfusion in the preconditioned livers. In a murine model of partial hepatic ischemia, the same group showed that IPC also inhibits apoptosis of hepatocytes at later stages of reperfusion [27].

The steatotic liver is particularly susceptible to ischemia-reperfusion injury resulting in poor outcome following liver surgery [44] and transplantation [45, 46]. There is, therefore, an urgent need for strategies against ischemia-reperfusion injury to increase the number of organs

available for liver transplantation and, moreover, improve the outcomes after transplantation of fatty livers and after hepatic resections. The recent report by Serafini et al. [47] shows IPC increases the tolerance of fatty livers to ischemia-reperfusion injury in rats. In this study obese Zucker rats subjected to 60 min of lobar liver ischemia had 70% survivors at 30 days in the ischemically preconditioned group as compared to no survivors in the ischemic controls.

Studies on Cold Ischemia

The protective effect of IPC is not restricted to warm ischemia and decreased tissue damage in cold preserved livers (cold storage-reperfusion injury) after IPC has been demonstrated in small and large animal models. In rat livers, IPC prior to storage in cold University of Wisconsin (UW) solution for 30 h decreased SEC death and Kupffer cell (KC) activation [48, 49]. In another study combining two sets of experiments, IPC prior to preservation of rat livers in cold UW solution for 30 h decreased SEC detachment and activities of matrix metalloproteinases, and also decreased SEC apoptosis after 1 h of reperfusion in an isolated perfused rat liver model [50]. In a rat liver transplant model IPC protected liver grafts from ischemia-reperfusion injury [26]. Furthermore, in a recent study in

cold-preserved rat livers Arai et al. [51] have observed that the benefit of IPC extends not only to the ipsilateral lobe, but also to the contralateral lobe resulting in an improved graft survival after orthotopic liver transplantation. In this study the authors observe that 'such heterologous preconditioning provides a new means to protect liver tissue against ischemia reperfusion injury without imposing ischemia on the target tissue' [51]. IPC also increased resistance to cold ischemic liver injury in pigs [52, 53].

Studies on Isolated Hepatocytes

The hepatoprotective response of IPC has also been shown in isolated hepatocytes. In *in vitro* studies on freshly isolated hepatocytes, preconditioned hepatocytes showed increased resistance to cell killing during hypoxic incubation [54, 55]. Another study has shown IPC-improved hepatocyte viability and energy metabolism in a model of isolated rat hepatocytes subjected to hypothermic preservation injury preceded by normothermic ischemia [56].

The above studies mostly in rodent livers have shown liver protection by IPC to warm and cold ischemia. However, there are a few published studies which have suggested that hepatic IPC may have only limited benefit. A study by Adam et al. [57] in fact suggested that preconditioning had a deleterious effect on hepatic tolerance to cold ischemia. This study used a model of isolated perfused livers from Wistar rats [57]. Preconditioning protocol of 5 or 10 min ischemia followed by 10 min reperfusion before liver harvesting, prior to extended cold ischemia of 24 h resulted in extensive reperfusion injury, increased vascular resistance and increased transaminases and LDH release. In a larger animal model using pigs, a preconditioning protocol of repeated 10 min ischemia followed by 10 min reperfusion, prior to 120 min or 200 min sustained ischemia was tested [58]. In the 120-min ischemia group IPC increased bile flow and ATP, but the degree of necrosis and apoptosis was not different from the control group. With 200 min ischemia IPC resulted in no significant differences in bile flow, ATP and liver enzymes from control group, and the degree of necrosis and apoptosis was in fact greater in preconditioned livers. This study suggested that IPC conferred some functional protection against reversible ischemia but no protection from prolonged ischemia in pigs [58]. The major difference between this study and those showing benefits with IPC is the use of three cycles of preconditioning in comparison with a single episode. In a more recent study Rudiger et al. [36] noted that in mice IPC

resulted in 100% animal survival with no morphological parenchymal injury after 75 min sustained ischemia as against 14% survival with significant parenchymal injury after 120 min ischemia.

Thus, a large body of evidence favors liver protection by IPC from injury in both warm and cold ischemia. The existence of IPC in the liver has been demonstrated in rodents, pig and humans. Although most of the data on hepatic IPC has been gathered in rodents and it is recognized that information on preconditioning in rats may not always be extrapolated to larger species and humans, the recent report by Clavien et al. [43] of the first human study is a thoughtful example of potential clinical application of the preconditioning effect.

Possible Mechanisms of Preconditioning

The precise mechanism of the IPC response is unknown. From studies on preconditioned myocardium, it is widely accepted that IPC is mediated via a receptor-targeting mechanism [59, 60]. Molecules released during ischemia attach to cellular receptors and contribute to preconditioning response. The candidate compounds implicated in liver IPC include adenosine [17, 20, 25, 48, 49], protein kinase C (PKC) [53–55], nitric oxide (NO) [13, 17, 20, 26, 41, 42], heat shock proteins (HSPs) [29, 61], tyrosine kinases [52], mitogen-activated protein kinases [55], oxidative stress [35, 50], nuclear factor κ B (NF- κ B) [62, 63], and modulation of apoptosis cascade [14, 27]. However the characterizations of these candidate compounds into different processes in the preconditioning cascade such as initiating trigger, signalling pathway and end effector are not defined and the interrelationship between these processes is unknown. In the liver, the most investigated molecules are NO [13], adenosine [49], PKC [54] and HSPs [29]. This article reviews the major developments in the characterization of these proposed mechanisms of preconditioning (fig. 2). Other mechanisms will not be discussed in further detail.

The Role of Adenosine

Adenosine is an extracellular molecule proposed both as 'trigger' and 'mediator' of IPC [3]. During ischemia, adenosine triphosphate is degraded to adenosine. The extracellular adenosine released in large quantities during ischemia is believed to play a role in the protective effect of IPC during reperfusion of ischemic tissue. Ischemia-reperfusion injury is associated with neutrophil and leukocyte activation and primary microvascular failure.




Fig. 2. Schematic illustration of possible mechanisms involved in IPC. NOS = Nitric oxide synthase; KATP = potassium-dependent ATP channel; cGMP = cyclic guanosine monophosphate; MAPK = mitogen-activated protein kinase.

Adenosine inhibits leukocyte adhesion, decreases expression of adhesion molecules and inhibits neutrophil and platelet function [64, 65]. Adenosine also inhibits free radical production [66, 67], which are important mediators of cellular damage in the early phase of ischemia-reperfusion injury, and is a potent vasodilator [68]. The

above would suggest adenosine may be protective against ischemia-reperfusion injury and the effects of adenosine in IPC are likely to be multifactorial. Most of the data on the role of adenosine in IPC has been gathered in cardiac muscle [69–71] and extrapolated to skeletal muscle [5] and kidneys [8].

Over the recent years a few studies have gathered evidence of the involvement of adenosine in liver IPC. Whereas A1 receptors have been implicated in the myocardium [72], A2 receptors have been proposed to be the adenosine receptor subtype likely to be expressed in the liver [49]. The existence of adenosine A2 receptors on hepatic SEC is supported indirectly by demonstrating a dose-dependent increase in cAMP by adenosine and selective A2 receptor agonist CGS-21680 [49]. In this study by Arai et al. [49], adenosine A2 receptor blockade prevented the protective effect of IPC in rat livers preserved in cold UW solution. IPC and administration of adenosine A2 receptor agonist, in this study, decreased SEC death and increased cAMP level [49]. The authors have proposed that SEC protection by IPC is mediated by activation of adenosine A2 receptors producing an increase in cAMP levels in SEC, but the mechanism downstream to increased cAMP, by which adenosine decreases SEC injury, is not explained. The same authors have previously shown that IPC suppressed KC activation and have stipulated the involvement of adenosine A2 receptors in this response [48]. IPC-induced protection of SECs can have profound implications for preservation of livers for transplantation, since SECs are more susceptible to cold preservation injury [73, 74] in contrast to hepatocytes which are vulnerable to warm ischemia-reperfusion injury [75]. SEC injury rather than hepatocellular injury has been shown to be responsible for graft failure from cold ischemia-reperfusion injury [73, 74, 76]. Peralta et al. [17] have postulated activation of adenosine A2 receptors with subsequent formation of NO to play a role in mediating IPC against warm ischemia-reperfusion injury. In this study adenosine administration in the presence of a NO donor reproduced the protective effect of IPC on hepatic parenchymal cells. In another study, a 3-fold increase in adenosine after IPC was associated with decreased parenchymal tissue damage [20]. Both IPC and increasing endogenous adenosine concentrations with the adenosine uptake inhibitor dipyridamole decreased hepatic leukocyte/endothelial cell interactions after ischemia-reperfusion injury [25]. All of the above studies have been carried out in rats and although the evidence is limited, suggest that adenosine modulates IPC-induced protection of non-parenchymal and parenchymal cells against cold and warm hepatic ischemia-reperfusion injury in the rat liver. There are no studies challenging the involvement of adenosine in the rat liver.

The data on adenosine from IPC studies in rat livers contradicts the information gathered in the rat heart. The role of adenosine in myocardial preconditioning is sup-

ported indirectly by studies in rabbits [77], pigs [78], dogs [79], and humans [80] demonstrating abolition of preconditioning by adenosine receptor blockade. However in rats, it is evident that adenosine has no role in IPC of the myocardium [81]. IPC is effective in the absence of extracellular accumulation of adenosine in the rat heart [82]. Thus, adenosine does not appear to be an endogenous trigger or obligatory mediator of preconditioning in rat hearts. Thus, in the rat species the adenosine concept does not seem to apply consistently to different tissues. It therefore seems likely that adenosine may only be a mediator to IPC of the liver but not a sole mechanism.

The Role of PKC

The PKC-mediated signalling pathway of myocardial preconditioning was proposed by Downey and colleagues [60, 83]. The hypothesis proposes that during preconditioning ischemia G protein activation following G protein coupling with adenosine receptors leads to PKC activation and subsequent translocation from the cytosol to the membrane where it phosphorylates substrate proteins to induce tolerance to subsequent ischemia [60]. However conflicting results in some species, particularly large animals where the concept does not apply consistently, would suggest that PKC activation is an epiphenomenon or secondary effect and not a primary mediator of the cardioprotective effects of preconditioning [84, 85]. Most of the evidence surrounding the PKC hypothesis is indirect and based on a pharmacological approach using PKC activators and inhibitors. Many of the inhibitors are not specific to PKC and are also isoform nonspecific. The above reviewers [84, 85] highlighted the limitations of pharmacological methods and also the fact that studies using isoform specific antibodies may not indicate activity of these specific PKC isoforms. Further, information on events downstream of PKC activation and the end effector of preconditioning is lacking at present.

In recent years, few studies have evaluated the evidence for involvement of PKC in preconditioning of the liver. This evidence is indirect and based on a pharmacological approach. Carini et al. [54] used an in vitro model of isolated rat hepatocytes and proposed that hypoxic preconditioning was mediated via PKC-mediated activation of vacuolar proton ATPase. In this study the increased tolerance of preconditioned hepatocytes to hypoxia was abolished by inhibition of PKC with chelerythrine or blocking vacuolar proton ATPase with bafilomycin A1 and mimicked by stimulators of PKC, 4 β -phorbol-12-myristate-13-acetate (PMA) and 1,2-dioctanoyl-glycerol (1,2-DOG). The authors observed that the prevention of

intracellular acidosis and of cytosolic Na⁺ increase during hypoxia was associated with decreased hypoxic injury in preconditioned hepatocytes [54]. In another study, the same authors observed that preconditioning was abolished by adenosine A2a receptor antagonist and have proposed a signalling pathway involving adenosine A2a receptors, PKC and kinases downstream of PKC (p38 mitogen-activated protein kinase) to be involved in hypoxic preconditioning of isolated rat hepatocytes [55]. However downstream of this point, the mechanisms by which liver injury is decreased have not been explained. In the heart, it has been suggested that the kinase cascade activated during preconditioning leads to the opening of mitochondrial K_{ATP} channels [86] but there is no evidence that these are the end effectors. There is data to suggest that mitochondrial K_{ATP} channels may simply act as another signal transduction step [87]. The kinase cascade can also stimulate phosphorylation of HSPs [88], activation of the transcription factor NF- κ B [89] and upregulation of inducible NO synthase [90] but the link with end effects of preconditioning has not been established. Ricciardi et al. [52, 53] have extended support for involvement of PKC and tyrosine kinase in liver IPC in larger animals. In one study, tolerance of ischemically preconditioned pig livers to cold ischemia was abolished by pretreatment with the PKC inhibitor chelerythrine [53]. In another study by the same authors pretreatment with tyrosine the kinase inhibitor genistein abolished the preconditioning effect in cold-preserved pig livers [52]. While these data support a role for PKC in IPC, they still do not prove that PKC is responsible for preconditioning.

The Role of HSP

HSPs are intracellular stress proteins that have been shown to accumulate after hyperthermia and ischemia [91]. The concept of sublethal whole animal hyperthermia conferring tolerance to other stresses such as ischemia and lethal endotoxin exposure is referred to as hyperthermic preconditioning and has been associated with HSP accumulation [92, 93]. In the rat liver, tolerance to ischemic injury has been associated with production of various inducible HSPs: HSP72 [30, 94], HSP73 [30] and HSP70 and HO-1/HSP32 [95, 96]. Ishikawa et al. [61] have proposed that in heat shock-preconditioned rat livers HSPs maintain mitochondrial membrane integrity during the ischemic episode, to produce energy-rich phosphates during reperfusion and thus contribute to ischemic tolerance. In an in vivo study in rats by Kume et al. [29] the reduced postischemic hepatocellular injury and improved survival was associated with overexpression of HSP72 in ischemi-

cally preconditioned livers as well as in the livers preconditioned with heat shock. In this study HSP72 was detected within 6–72 h after heat exposure and the authors have proposed that HSP72 production is associated with a delayed protective effect of IPC. The link between HSP72 and delayed effect of IPC has not been explained. It is also not clear whether HSP production and accumulation is the reason for resistance to ischemia or merely a marker of tolerance [97].

While these studies demonstrate that HSPs are detected after preconditioning, the molecular mechanism of protection associated with HSP accumulation is not explained and these studies do not prove that HSPs are responsible for preconditioning.

The Role of NO

NO is a colorless, odorless, free radical gas which has been identified as an important signaling molecule in almost every tissue in the body. NO is produced from L-arginine by the enzyme NO synthase. In the liver, as in many other organs NO has many actions and cellular sources. Recent evidence supports the role of NO in regulating perfusion of the hepatic microcirculation [98]. The breakdown of microvascular perfusion with subsequent impairment of tissue oxygenation plays a central role in the pathophysiology of ischemia-reperfusion-induced injury of the liver [99]. Treatment of rats with nonspecific NO synthase inhibitors resulted in a failure of microvascular perfusion and development of patchy necrosis [100, 101]. Augmentation of NO synthesis with NO donors has been shown to attenuate hepatic ischemia-reperfusion injury and improve posttransplant survival [102]. NO may modulate microvascular perfusion through its vasodilatory effect [103] and through its anti-inflammatory actions including inhibitory effects on stellate cell activation [104], neutrophil adhesion [105] and platelet aggregation [101].

It has been proposed that NO plays a key role in both initiating and mediating IPC. While functional evidence in the heart indicates that NO modulates both acute and delayed preconditioning, downstream of this point in the biochemical pathway hypotheses are less well established [106, 107]. A recent study by Lochner et al. [108] has proposed that NO through generation of cGMP acts as a trigger of acute preconditioning in rat hearts. Parratt [109] has suggested that endocardial endothelium-derived mediators such as NO may mediate cardioprotective response of IPC by elevation of cGMP, which in turn could reduce energy demand by limiting myocardial cAMP levels by stimulation of cGMP-sensitive cAMP phosphodies-

terase enzyme. It appears that whereas the acute phase of preconditioning is protein synthesis independent, the late phase requires new protein synthesis. It has been proposed that eNOS-derived NO leads to activation of PKC and other kinases, which in turn through NF- κ B and other transcription factors leads to an increase in transcription of iNOS [107]. The end effector of IPC in the supposed NO pathway is speculative and cGMP-dependent mechanisms and ATP-sensitive potassium channel have been proposed [107].

In the liver it has been suggested that depending on the rate of its production, NO may also play a mediating role in preconditioning [110]. NO has been implicated in IPC-associated decreased tissue damage in both warm ischemia [13] and cold ischemic storage [26] of the rat liver. However the link between protective effects of IPC and NO is speculative. Peralta et al. [13] suggested that liver IPC in rats is mediated by the inhibitory action of NO on endothelin. In other studies in rats, the same authors have demonstrated that inhibition of adenosine and simultaneous administration of NO donor offered similar results to IPC [17] and have proposed that activation of adenosine A2 receptors with subsequent NO formation mediates IPC in the rat liver [20]. Yin et al. [26] have postulated that IPC increased resistance to cold ischemic liver injury in rats through stimulation of endogenous NO. In this study pharmacological NO stimulation mimicked and NO inhibition antagonized IPC-associated protection of liver grafts from preservation reperfusion injury in a rat liver transplantation model but the mechanism has not been explained. Recently, our group has shown an increased hepatic intracellular oxygenation [42] and increased hepatic microcirculation [41] with IPC which was associated with increased NO levels. A recent report by Serafin et al. [47] has implicated NO in the preconditioning response for ischemia-reperfusion injury in fatty livers.

NO suppresses apoptosis in endothelial cells. In recent years it has been suggested that apoptosis is the dominant mechanism for cell turnover in the human liver. Apoptosis is a rapid process terminating in nuclear pyknosis and cell death. Recent evidence has shown that apoptosis of SEC and hepatocytes are a feature of ischemia-reperfusion injury in warm [111] and cold [112] ischemia of the liver. The signalling pathways leading to nuclear apoptosis in response to extracellular stimuli involve activation of cysteine proteases known as caspases and release of cytochrome c from the mitochondria [113]. Subsequent activation of downstream caspases such as caspase 3 ultimately executes nuclear apoptosis [114]. Antiapoptotic molecules such as Bcl-2 and caspase inhibitors have been

shown to prevent release of mitochondrial cytochrome c [115]. In an experimental model of partial hepatic ischemia, IPC inhibited apoptosis of SEC and hepatocytes and was associated with inhibition of caspase 3 activity [27]. In the study IPC was not associated with higher Bcl-2 or Bcl-xl expression. The link between IPC and inhibition of caspase activity is speculative. NO has been shown to inhibit caspase activity in vitro [116]. Apoptosis in hepatocytes exposed to TNF- α and actinomycin D was prevented by NO. In this study NO produced by an NO donor or through iNOS gene expression inhibited caspase family proteases by S-nitrosylation and prevented cytochrome c release [113]. Other mechanisms for an antiapoptotic effect of NO are increase in cGMP [116] and upregulation of Bcl-2 [117] and HSPs [118]. Thus, potentially liver IPC may be mediated through NO modulation of apoptosis cascade.

Role of IPC in Hepatic Surgery

Ischemia-reperfusion injury is a major cause of morbidity and mortality following liver surgery and transplantation. In the setting of liver resections, the effects of intermittent inflow occlusion, continuous inflow occlusion and total vascular exclusion during liver resections have been studied in clinical trials [119–121]. Whereas total vascular exclusion was effective in reducing blood loss, it led to unpredictable hemodynamic intolerance, increased morbidity and longer hospital stay [119]. This is not surprising since the state of total vascular exclusion is akin to the anhepatic phase of liver transplantation and hemodynamic consequences on reperfusion would be anticipated. In a prospective evaluation of intermittent inflow occlusion versus no inflow occlusion in patients undergoing liver resections, the former resulted in less blood loss and better preservation of liver function in the early postoperative period [120]. When intermittent versus continuous inflow occlusion were studied in patients undergoing liver resections [121], the group subjected to intermittent inflow occlusion was associated with decreased hepatocellular injury indicated by lower postoperative liver enzymes and serum bilirubin levels. However the intraoperative blood loss during liver transection was significantly higher in this group and this is most likely related to bleeding from the transected surface during successive reperfusion episodes [121]. Thus, the increased blood loss and likely increased duration of surgery due to successive reperfusion episodes may outweigh the benefit of intermittent occlusion on parenchymal tolerance to

ischemia. Although some liver resections can be performed without vascular inflow occlusion, prolonged ischemia may be unavoidable to achieve radical tumor resection. An ideal protective strategy for human liver surgery would allow a bloodless parenchymal transection and an increased parenchymal tolerance to ischemia. In theory, IPC may obviate the need for intermittent releases of hepatic vascular occlusions and extend safe periods of ischemia by increasing hepatic tolerance to ischemia during hepatic surgery. The potential for clinical application of IPC for hemihepatectomies under inflow occlusion has been demonstrated by Clavien et al. [43]. In this study IPC protected against 30 min of inflow occlusion with patients showing a 2-fold decrease in serum transaminases compared to patients subjected to continuous ischemia only, but no significant differences in duration of surgery, need for intensive care or mortality. This study provides evidence that IPC occurs in the human liver.

In the setting of liver transplantation, ischemia time of the donor liver is a major determinant of graft outcome and patient survival after liver transplantation [122]. Liver transplantation requires mandatory organ ischemia. Warm ischemia to the graft may occur at organ harvest in an unstable donor and cold ischemia occurs during preservation of the liver for transplantation. During implantation of the graft in the recipient, the liver is subjected to further warm ischemia until the vascular anastomoses are completed. Finally reperfusion injury is inevitable following revascularization. Prolonged ischemia results in primary nonfunction or dysfunction of the transplanted liver graft and is associated with biliary and vascular complications [122, 123] often resulting in retransplantation. This adversely affects patient outcome and survival. Therefore

IPC is an attractive strategy to assist liver preservation and protect the liver from ischemia-reperfusion injury during transplantation by increasing ischemic tissue tolerance of the liver. As yet there are no reported studies demonstrating clinical benefits of IPC in patients undergoing liver transplantation. Most animal studies have shown that IPC offers a degree of protection against cold ischemia in experimental liver transplantation. This data in animal models is encouraging and clinical studies are required to clarify the potential application of IPC in human liver transplantation.

Conclusions

The past decade has provided interesting new data establishing the existence of IPC in the liver. IPC is a powerful endogenous means to protect the liver from ischemia. To date one study has demonstrated human clinical benefits of liver IPC. Further clinical studies are required to prove unequivocally that IPC is possible in the human liver. However the central mechanism of IPC remains undefined. Current research has demonstrated that IPC is an endogenous adaptive phenomenon that can be reproduced easily in different models of warm and cold ischemia, and in animals as well as humans. However the causal relationship between the initiating event, biochemical pathways and end effector molecules remains mechanistically undefined and controversial. As the field advances with mechanistically descriptive studies, these controversies in interrelationships in the preconditioning cascade are likely to be resolved and will lead to pharmacological strategies for protecting the liver from ischemic injury.

References

- 1 Bilzer M, Gerbes AL: Preservation injury of the liver: Mechanisms and novel therapeutic strategies. *J Hepatol* 2000;32:508–515.
- 2 Murry CE, Jennings RB, Reimer KA: Preconditioning with ischemia: A delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;74:1124–1136.
- 3 Downey JM, Liu GS, Thornton JD: Adenosine and the anti-infarct effects of preconditioning. *Cardiovasc Res* 1993;27:3–8.
- 4 Yellon DM, Alkhulaifi AM, Pugsley WB: Preconditioning of the human myocardium. *Lancet* 1993;343:276–277.
- 5 Pang CY, Yang RZ, Zhong A, Xu N, Boyd B, Forrest CR: Acute ischaemic preconditioning protects against skeletal muscle infarction in the pig. *Cardiovasc Res* 1995;29:782–788.
- 6 Heurteaux C, Lauritzen I, Widmann C, Lazdunski M: Essential role of adenosine, adenosine A1 receptors, and ATP-sensitive K⁺ channels in cerebral ischemic preconditioning. *Proc Natl Acad Sci USA* 1995;92:4666–4670.
- 7 Matsuyama K, Chiba Y, Ihaya A, Kimura T, Tanigawa N, Muraoka R: Effect of spinal cord preconditioning on paraplegia during cross-clamping of the thoracic aorta. *Ann Thorac Surg* 1997;63:1315–1320.
- 8 Turman MA, Bates CM: Susceptibility of human proximal tubular cells to hypoxia: Effect of hypoxic preconditioning and comparison to glomerular cells. *Ren Fail* 1997;19:47–60.
- 9 Hotter G, Closa D, Prados M, et al: Intestinal preconditioning is mediated by a transient increase in nitric oxide. *Biochem Biophys Res Commun* 1996;222:27–32.
- 10 Lloris-Carsi JM, Cejalvo D, Toledo-Pereyra LH, Calvo MA, Suzuki S: Preconditioning: Effect upon lesion modulation in warm liver ischemia. *Transplant Proc* 1993;25:3303–3304.
- 11 Hardy KJ, McClure DN, Subwongcharoen S: Ischaemic preconditioning of the liver: A preliminary study. *Aust NZ J Surg* 1996;66:707–710.
- 12 Yoshizumi T, Yanaga K, Soejima Y, Maeda T, Uchiyama H, Sugimachi K: Amelioration of liver injury by ischaemic preconditioning. *Br J Surg* 1998;85:1636–1640.

- 13 Peralta C, Closa D, Hotter G, Gelpi E, Prats N, Rosello-Catafau J: Liver ischemic preconditioning is mediated by the inhibitory action of nitric oxide on endothelin. *Biochem Biophys Res Commun* 1996;229:264-270.
- 14 Peralta C, Perales JC, Bartrons R, et al: The combination of ischemic preconditioning and liver Bcl-2 overexpression is a suitable strategy to prevent liver and lung damage after hepatic ischemia-reperfusion. *Am J Pathol* 2002;160:2111-2122.
- 15 Teoh N, Dela Pena A, Farrell G: Hepatic ischemic preconditioning in mice is associated with activation of NF-kappaB, p38 kinase, and cell cycle entry. *Hepatology* 2002;36:94-102.
- 16 Fernandez L, Heredia N, Grande L, et al: Preconditioning protects liver and lung damage in rat liver transplantation: Role of xanthine/xanthine oxidase. *Hepatology* 2002;36:562-572.
- 17 Peralta C, Hotter G, Closa D, Gelpi E, Bulbena O, Rosello-Catafau J: Protective effect of preconditioning on the injury associated to hepatic ischemia-reperfusion in the rat: Role of nitric oxide and adenosine. *Hepatology* 1997;25:934-937.
- 18 Peralta C, Closa D, Xaus C, Gelpi E, Rosello-Catafau J, Hotter G: Hepatic preconditioning in rats is defined by a balance of adenosine and xanthine. *Hepatology* 1998;28:768-773.
- 19 Peralta C, Prats N, Xaus C, Gelpi E, Rosello-Catafau J: Protective effect of liver ischemic preconditioning on liver and lung injury induced by hepatic ischemia-reperfusion in the rat. *Hepatology* 1999;30:1481-1489.
- 20 Peralta C, Hotter G, Closa D, et al: The protective role of adenosine in inducing nitric oxide synthesis in rat liver ischemia preconditioning is mediated by activation of adenosine A2 receptors. *Hepatology* 1999;29:126-132.
- 21 Peralta C, Bartrons R, Riera L, et al: Hepatic preconditioning preserves energy metabolism during sustained ischemia. *Am J Physiol Gastrointest Liver Physiol* 2000;279:G163-G171.
- 22 Peralta C, Bartrons R, Serafin A, et al: Adenosine monophosphate-activated protein kinase mediates the protective effects of ischemic preconditioning on hepatic ischemia-reperfusion injury in the rat. *Hepatology* 2001;34:1164-1173.
- 23 Peralta C, Fernandez L, Panes J, et al: Preconditioning protects against systemic disorders associated with hepatic ischemia-reperfusion through blockade of tumor necrosis factor-induced P-selectin up-regulation in the rat. *Hepatology* 2001;33:100-113.
- 24 Nilsson B, Friman S, Gustafsson BI, Delbro DS: Preconditioning protects against ischemia/reperfusion injury of the liver. *J Gastrointest Surg* 2000;4:44-49.
- 25 Howell JG, Zibari GB, Brown MF, et al: Both ischemic and pharmacological preconditioning decrease hepatic leukocyte/endothelial cell interactions. *Transplantation* 2000;69:300-303.
- 26 Yin DP, Sankary HN, Chong AS, et al: Protective effect of ischemic preconditioning on liver preservation-reperfusion injury in rats. *Transplantation* 1998;66:152-157.
- 27 Yadav SS, Sindram D, Perry DK, Clavien PA: Ischemic preconditioning protects the mouse liver by inhibition of apoptosis through a caspase-dependent pathway. *Hepatology* 1999;30:1223-1231.
- 28 Nakayama H, Yamamoto Y, Kume M, et al: Pharmacologic stimulation of adenosine A2 receptor supplants ischemic preconditioning in providing ischemic tolerance in rat livers. *Surgery* 1999;126:945-954.
- 29 Kume M, Yamamoto Y, Saad S, et al: Ischemic preconditioning of the liver in rats: Implications of heat shock protein induction to increase tolerance of ischemia-reperfusion injury. *J Lab Clin Med* 1996;128:251-258.
- 30 Kume M, Yamamoto Y, Yamagami K, Ishikawa Y, Uchinami H, Yamaoka Y: Pharmacological hepatic preconditioning: Involvement of 70-kDa heat shock proteins (HSP72 and HSP73) in ischaemic tolerance after intravenous administration of doxorubicin. *Br J Surg* 2000;87:1168-1175.
- 31 Tsuyama H, Shimizu K, Yoshimoto K, et al: Protective effect of ischemic preconditioning on hepatic ischemia-reperfusion injury in mice. *Transplant Proc* 2000;32:2310-2313.
- 32 Yamada F, Abe T, Saito T, Tsuciya T, Ishii S, Gotoh M: Ischemic preconditioning enhances regenerative capacity of hepatocytes after prolonged ischemia. *Transplant Proc* 2001;33:956.
- 33 Saito T, Ishii S, Abe T, et al: Effect of preconditioning in the liver against ischemia/reperfusion injury, protection of sinusoidal cells and alterations of gene transcription. *Transplant Proc* 2001;33:849.
- 34 Zhang Y, Zhang B, Pan R: Protective effect of ischemic preconditioning on liver. *Chin J Traumatol* 2001;4:123-125.
- 35 Peralta C, Bulbena O, Xaus C, et al: Ischemic preconditioning: A defense mechanism against the reactive oxygen species generated after hepatic ischemia reperfusion. *Transplantation* 2002;73:1203-1211.
- 36 Rudiger HA, Kang K-J, Sindram D, Riehle H-M, Clavien PA: Comparison of ischemic preconditioning and intermittent and continuous inflow occlusion in the murine liver. *Ann Surg* 2002;235:400-407.
- 37 Ajamieh H, Merino N, Candelario-Jalil E, et al: Similar protective effect of ischaemic and ozone oxidative preconditionings in liver ischaemia/reperfusion injury. *Pharmacol Res* 2002;45:333-339.
- 38 Iwasaki Y, Tagaya N, Hattori Y, Yamaguchi K, Kubota K: Protective effect of ischemic preconditioning against intermittent warm-ischemia-induced liver injury. *J Surg Res* 2002;107:82-92.
- 39 Ishii S, Abe T, Saito T, et al: Effects of preconditioning on ischemia/reperfusion injury of hepatocytes determined by immediate early gene transcription. *J Hepatobiliary Pancreat Surg* 2001;8:461-468.
- 40 Zapletal C, Maksan SM, Lehmann T, et al: Ischemic preconditioning improves liver microcirculation after ischemia/reperfusion. *Transplant Proc* 1999;31:3260-3262.
- 41 Koti RS, Yang W, Dashwood MR, Davidson BR, Seifalian AM: Effect of ischemic preconditioning on hepatic microcirculation and function in a rat model of ischemia reperfusion injury. *Liver Transpl* 2002;8:1182-1191.
- 42 Koti RS, Yang W, McBride AG, Seifalian AM, Davidson BR: The relationship of hepatic tissue oxygenation with nitric oxide metabolism in ischemic preconditioning of the liver. *FASEB J* 2002;16:1654-1656.
- 43 Clavien PA, Yadav S, Sindram D, Bentley RC: Protective effects of ischemic preconditioning for liver resection performed under inflow occlusion in humans. *Ann Surg* 2000;232:155-162.
- 44 Makuuchi M, Mori T, Gunven P, et al: Safety of hemihepatic vascular occlusion during resection of the liver. *Surg Gynecol Obstet* 1989;130:824-831.
- 45 Todo S, Demetris AJ, Makowka L, et al: Primary nonfunction of hepatic allografts with preexisting fatty infiltration. *Transplantation* 1989;47:903.
- 46 Adam R, Reynes M, Johann M, et al: The outcome of steatotic grafts in liver transplantation. *Transplant Proc* 1991;23:1538-1540.
- 47 Serafin A, Rosello-Catafau J, Prats N, Xaus C, Gelpi E, Peralta C: Ischemic preconditioning increases the tolerance of fatty liver to hepatic ischemia-reperfusion injury in the rat. *Am J Pathol* 2002;161:587-601.
- 48 Arai M, Thurman RG, Lemasters JJ: Involvement of Kupffer cells and sinusoidal endothelial cells in ischemic preconditioning to rat livers stored for transplantation. *Transplant Proc* 1999;31:425-427.
- 49 Arai M, Thurman RG, Lemasters JJ: Contribution of adenosine A(2) receptors and cyclic adenosine monophosphate to protective ischemic preconditioning of sinusoidal endothelial cells against storage/reperfusion injury in rat livers. *Hepatology* 2000;32:297-302.
- 50 Sindram D, Rudiger HA, Upadhyya AG, Strasberg SM, Clavien PA: Ischemic preconditioning protects against cold ischemic injury through an oxidative stress dependent mechanism. *J Hepatol* 2002;36:78-84.
- 51 Arai M, Thurman RG, Lemasters JJ: Ischemic preconditioning of rat livers against cold storage-reperfusion injury: Role of nonparenchymal cells and the phenomenon of heterologous preconditioning. *Liver Transpl* 2001;7:292-299.
- 52 Ricciardi R, Schaffer BK, Kim RD, et al: Protective effects of ischemic preconditioning on the cold-preserved liver are tyrosine kinase dependent. *Transplantation* 2001;72:406-412.
- 53 Ricciardi R, Meyers WC, Schaffer BK, et al: Protein kinase C inhibition abrogates hepatic ischemic preconditioning responses. *J Surg Res* 2001;97:144-149.
- 54 Carini R, De Cesaris MG, Splendore R, Bagnati M, Albano E: Ischemic preconditioning reduces Na(+) accumulation and cell killing in isolated rat hepatocytes exposed to hypoxia. *Hepatology* 2000;31:166-172.

- 55 Carini R, Cesaris MG, Splendore R, et al: Signal pathway involved in the development of hypoxic preconditioning in rat hepatocytes. *Hepatology* 2001;33:131–139.
- 56 Compagnon P, Wang H, Southard J, Mangino M: Ischemic preconditioning in a rodent hepatocyte model of liver hypothermic preservation injury. *Cryobiology* 2002;44:269–278.
- 57 Adam R, Arnault I, Bao YM, Salvucci M, Sebah M, Bismuth H: Effect of ischemic preconditioning on hepatic tolerance to cold ischemia in the rat. *Transpl Int* 1998;11(suppl 1):S168–S170.
- 58 Schulz R, Walz MK, Behrends M, Neumann T, Gerken G, Heusch G: Minimal protection of the liver by ischemic preconditioning in pigs. *Am J Physiol Heart Circ Physiol* 2001;280:H198–H207.
- 59 Dekker LR: Toward the heart of ischemic preconditioning. *Cardiovasc Res* 1998;37:14–20.
- 60 Nakano A, Cohen MV, Downey JM: Ischemic preconditioning. From basic mechanisms to clinical applications. *Pharmacol Ther* 2000;86:263–275.
- 61 Ishikawa Y, Yamamoto Y, Kume M, et al: Heat shock preconditioning on mitochondria during warm ischemia in rat livers. *J Surg Res* 1999;87:178–184.
- 62 Funaki H, Shimizu K, Harada S, et al: Essential role for nuclear factor kappaB in ischemic preconditioning for ischemia-reperfusion injury of the mouse liver. *Transplantation* 2002;74:551–556.
- 63 Ricciardi R, Shah SA, Wheeler SM, et al: Regulation of NFkappaB in hepatic ischemic preconditioning. *J Am Coll Surg* 2002;195:319–326.
- 64 Bouma MG, van den Wildenberg FA, Buurman WA: The anti-inflammatory potential of adenosine in ischemia-reperfusion injury: Established and putative beneficial actions of a retaliatory metabolite. *Shock* 1997;8:313–320.
- 65 Keller MW, Geddes L, Spotnitz W, Kaul S, Duling BR: Microcirculatory dysfunction following perfusion with hyperkalemic, hypothermic, cardioplegic solutions and blood reperfusion. Effects of adenosine. *Circulation* 1991;84:2485–2494.
- 66 Nolte D, Lehr HA, Messmer K: Adenosine inhibits postischemic leukocyte-endothelium interaction in postcapillary venules of the hamster. *Am J Physiol* 1991;261:H651–H655.
- 67 Cronstein BN: Adenosine, an endogenous anti-inflammatory agent. *J Appl Physiol* 1994;76:5–13.
- 68 Hori M, Kitakaze M: Adenosine, the heart, and coronary circulation. *Hypertension* 1991;18:565–574.
- 69 Downey JM, Liu GS, Thornton JD: Adenosine and the anti-infarct effects of preconditioning. *Cardiovasc Res* 1993;27:3–8.
- 70 Murry CE, Jennings RB, Reimer KA: New insights into potential mechanisms of ischemic preconditioning. *Circulation* 1991;84:442–445.
- 71 Lawson CS, Downey JM: Preconditioning: State of the art myocardial protection. *Cardiovasc Res* 1993;27:542–550.
- 72 Mullane K, Bullough D: Harnessing an endogenous cardioprotective mechanism: Cellular sources and sites of action of adenosine. *J Mol Cell Cardiol* 1995;27:1041–1054.
- 73 Caldwell-Kenkel JC, Thurman RG, Lemasters JJ: Selective loss of nonparenchymal cell viability after cold ischemic storage of rat livers. *Transplantation* 1988;45:834–837.
- 74 Caldwell-Kenkel JC, Currin RT, Tanaka Y, Thurman RG, Lemasters JJ: Reperfusion injury to endothelial cells following cold ischemic storage of rat livers. *Hepatology* 1989;10:292–299.
- 75 Noack K, Bronk SF, Kato A, Gores GJ: The greater vulnerability of bile duct cells to reoxygenation injury than to anoxia. Implications for the pathogenesis of biliary strictures after liver transplantation. *Transplantation* 1993;56:495–500.
- 76 Ikeda T, Yanaga K, Kishikawa K, Kakizoe S, Shimada M, Sugimachi K: Ischemic injury in liver transplantation: Difference in injury sites between warm and cold ischemia in rats. *Hepatology* 1992;16:454–461.
- 77 Liu GS, Thornton J, Van Winkle DM, Stanley AW, Olsson RA, Downey JM: Protection against infarction afforded by preconditioning is mediated by A1 adenosine receptors in rabbit heart. *Circulation* 1991;84:350–356.
- 78 Schulz R, Rose J, Post H, Heusch G: Involvement of endogenous adenosine in ischaemic preconditioning in swine. *Pflügers Arch* 1995;430:273–282.
- 79 Auchampach JA, Gross GJ: Adenosine A1 receptors and, K-ATP channels, and ischemic preconditioning in dogs. *Am J Physiol* 1993;264:H1327–H1336.
- 80 Walker DM, Walker JM, Pugsley WB, Pattison CW, Yellon DM: Preconditioning in isolated superfused human muscle. *J Mol Cell Cardiol* 1995;27:1349–1357.
- 81 Cave AC, Collis CS, Downey JM, Hearse DJ: Improved functional recovery by ischaemic preconditioning is not mediated by adenosine in the globally ischaemic isolated rat heart. *Cardiovasc Res* 1993;27:663–668.
- 82 Lasley RD, Anderson GM, Mentzer RM: Ischaemic and hypoxic preconditioning enhance postischaemic recovery of function in the rat heart. *Cardiovasc Res* 1993;27:565–570.
- 83 Liu Y, Ytrehus K, Downey JM: Evidence that translocation of protein kinase C is a key event during ischemic preconditioning of rabbit myocardium. *J Mol Cell Cardiol* 1994;26:661–668.
- 84 Brooks G, Hearse DJ: Role of protein kinase C in ischemic preconditioning: Player or spectator? *Circ Res* 1996;79:627–630.
- 85 Simkhovich BZ, Przyklenk K, Kloner RA: Role of protein kinase C as a cellular mediator of ischemic preconditioning: A critical review. *Cardiovasc Res* 1998;40:9–22.
- 86 Baines CP, Liu GS, Birincioglu M, Critz SD, Cohen MV, Downey JM: Ischemic preconditioning depends on interaction between mitochondrial KATP channels and actin cytoskeleton. *Am J Physiol* 1999;276:H1361–H1368.
- 87 Pain T, Cohen MV, Downey JM: The mitochondrial KATP channel may be a trigger rather than the end-effector of preconditioning's anti-infarct effect. *Circulation* 1999;100(suppl I):I–342.
- 88 Rouse J, Cohen P, Trigon S, et al: A novel kinase cascade triggered by stress and heat shock that stimulates MAPKAP kinase-2 and phosphorylation of the small heat shock proteins. *Cell* 1994;78:1027–1037.
- 89 Morgan EN, Boyle EM, Yun W, et al: An essential role for NF-kappa B in the cardioprotective response to ischemia. *Ann Thorac Surg* 1999;68:377.
- 90 Xuan YT, Tang XL, Qiu Y, et al: Biphasic response of cardiac NO synthase isoforms to ischemic preconditioning in conscious rabbits. *Am J Physiol* 2000;279:H2360.
- 91 Tissieres A, Mitchell HK, Tracy UM: Protein synthesis in salivary glands of *Drosophila melanogaster*: Relation to chromosome puffs. *J Mol Biol* 1974;84:389–398.
- 92 Ryan AJ, Flanagan SW, Moseley PL, Gisolfi CV: Acute heat stress protects rats against endotoxin shock. *J Appl Physiol* 1992;73:1517–1522.
- 93 Hotchkiss R, Nunnally I, Lindquist S, Taulien J, Perdrizet G, Karl I: Hyperthermia protects mice against the lethal effects of endotoxin. *Am J Physiol* 1993;265:R1447–R1457.
- 94 Saad S, Kanai M, Awane M, et al: Protective effect of heat shock pretreatment with heat shock protein induction before hepatic warm ischemic injury caused by Pringle's maneuver. *Surgery* 1995;118:510–516.
- 95 Terajima H, Enders G, Thiaener A, et al: Impact of hyperthermic preconditioning on postischemic hepatic microcirculatory disturbances in an isolated perfusion model of the rat liver. *Hepatology* 2000;31:407–415.
- 96 Terajima H, Thiaener A, Hammer C, Messmer K, Yamamoto Y, Yamaoka Y: Attenuation of hepatic microcirculatory failure during in situ xenogeneic rat liver perfusion by heat shock preconditioning. *Transplant Proc* 2000;32:1111.
- 97 Moseley PL: Heat shock proteins: A broader perspective. *J Lab Clin Med* 1996;128:233–234.
- 98 Clemens MG: Nitric oxide in liver injury. *Hepatology* 1999;30:1–5.
- 99 Koo A, Komatsu H, Tao G, Inoue M, Guth PH, Kaplowitz N: Contribution of no-reflow phenomenon to hepatic injury after ischemia-reperfusion: Evidence for a role for superoxide anion. *Hepatology* 1992;15:507–514.
- 100 Shibayama Y, Nakata K: Role of septal fibrosis in development of hepatic circulatory disturbance in the presence of liver cell enlargement. *Liver* 1992;12:84–89.
- 101 Harbrecht BG, Billiar TR, Stadler J, et al: Inhibition of nitric oxide synthesis during endotoxemia promotes intrahepatic thrombosis and an oxygen radical-mediated hepatic injury. *J Leukoc Biol* 1992;52:390–394.

- 102 Niu X-F, Smith CW, Kubes P: Intracellular oxidative stress induced by nitric oxide synthesis inhibition increased endothelial cell adhesivity for neutrophils. *Circ Res* 1994;74:1133-1140.
- 103 Jaeschke H, Smith CW, Clemens MG, Ganey PE, Roth RA: Mechanisms of inflammatory liver injury: Adhesion molecules and cytotoxicity of neutrophils. *Toxicol Appl Pharmacol* 1996;139:213-226.
- 104 Rockey DC, Chung JJ: Inducible nitric oxide synthase in rat hepatic lipocytes and the effect of nitric oxide on lipocyte contractility. *J Clin Invest* 1995;95:1199-1206.
- 105 Harbrecht BG, Wu B, Watkins SC, Billiar TR, Peitzman AB: Inhibition of nitric oxide synthesis during severe shock but not after resuscitation increases hepatic injury and neutrophil accumulation in hemorrhaged rats. *Shock* 1997;8:415-421.
- 106 Rakhit RD, Edwards JR, Marber MS: Nitric oxide, nitrates and ischaemic preconditioning. *Cardiovasc Res* 1999;43:621-627.
- 107 Nandagopal K, Dawson TM, Dawson VL: Critical role for nitric oxide signalling in cardiac and neuronal ischemic preconditioning and tolerance. *J Pharmacol Exp Ther* 2001;297:474-478.
- 108 Lochner A, Marais E, Genade S, Moolman JA: Nitric oxide: A trigger for classic preconditioning? *Am J Physiol Heart Circ Physiol* 2000;279:H2752-H2765.
- 109 Parratt J: Protection of the heart by ischaemic preconditioning: Mechanisms and possibilities for pharmacological exploitation. *Trends Pharmacol Sci* 1994;15:19-25.
- 110 Roth E: The impact of *L*-arginine-nitric oxide metabolism on ischemia/reperfusion injury. *Curr Opin Clin Nutr Metab Care* 1998;1:97-99.
- 111 Kohli V, Selzner M, Madden JF, Bentley RC, Clavien PA: Endothelial cell and hepatocyte deaths occur by apoptosis after ischemia-reperfusion injury in the rat liver. *Transplantation* 1999;67:1099-1105.
- 112 Gao W, Bentley RC, Madden JF, Clavien PA: Apoptosis of sinusoidal endothelial cells is a critical mechanism of preservation injury in rat liver transplantation. *Hepatology* 1998;27:1652-1660.
- 113 Kim Y-M, Kim T-H, Chung H-T, Talanian RV, Yin X-M, Billiar TR: Nitric oxide prevents tumor necrosis factor alpha-induced rat hepatocyte apoptosis by the interruption of mitochondrial apoptotic signaling through s-nitrosylation of caspase-8. *Hepatology* 2000;32:770-778.
- 114 Miller DG: The role of caspase family of cysteine proteases in apoptosis. *Semin Immunol* 1997;9:35-39.
- 115 Yang J, Liu X, Bhalla K, et al: Prevention of apoptosis by Bcl-2: Release of cytochrome c from mitochondria blocked. *Science* 1997;275:1129-1132.
- 116 Kim YM, Talanian RV, Billiar TR: Nitric oxide inhibits apoptosis by preventing increases in caspase-3-like activity via two distinct mechanisms. *J Biol Chem* 1997;272:31138-31148.
- 117 Genaro AM, Hortelano S, Alvarez A, Martinez C, Bosca L: Splenic B lymphocyte programmed cell death is prevented by nitric oxide release through mechanisms involving sustained Bcl-2 levels. *J Clin Invest* 1995;95:1884-1890.
- 118 Kim Y, de Vera ME, Watkins SC, Billiar TR: Nitric oxide protects cultured rat hepatocytes from tumor necrosis factor-alpha-induced apoptosis by inducing heat shock protein 70 expression. *J Biol Chem* 1997;272:1402-1411.
- 119 Belghiti J, Noun R, Zante E, Ballet T, Sauvanet A: Portal triad clamping or hepatic vascular exclusion for major liver resection: A controlled study. *Ann Surg* 1996;224:155-161.
- 120 Man K, Fan ST, Ng IO, Lo CM, Liu CL, Wong J: Prospective evaluation of Pringle maneuver in hepatectomy for liver tumors by a randomized study. *Ann Surg* 1997;226:704-713.
- 121 Belghiti J, Noun R, Malafosse R, et al: Continuous versus intermittent portal triad clamping for liver resection: A controlled study. *Ann Surg* 1999;229:369-375.
- 122 Lemasters JJ, Bunzendahl H, Thurman RG: Reperfusion injury to donor livers stored for transplantation. *Liver Transpl Surg* 1995;1:124-138.
- 123 Blankensteijn JD, Terpstra OT: Liver preservation: The past and the future. *Hepatology* 1991;13:1235-1250.